

Savey[®] *Ovicide / Miticide*

50-WP

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION

ACTIVE INGREDIENT:

% By Wt.

Hexythiazox:

trans-5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide.....50%

INERT INGREDIENTS50%

TOTAL 100%

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA-registered label.
- Do not apply this product through any type of irrigation system.

Crop	Pest	Rate oz/acre	Comments
Hops	Twospotted spider mite	4 - 6	Savey is active against all immature stages of spider mites, including freshly laid eggs. Apply Savey prior to adult mite buildup, up to burr formation in hop vines. In general, this period will coincide up to the time vines have reached a height of 18 feet or "to the wire". Complete coverage of leaf surfaces is essential. Apply by ground equipment in suitable volume and method to assure complete coverage of entire vines. Use higher rates for moderate to high spider mite egg infestations (10 – 15 eggs/leaf). If adult mites are present in moderate numbers (1 - 5 motile mites per leaf), apply Savey in combination with a mite adulticide.

- Use only one application per growing season.
- Apply Savey in conjunction with an integrated mite management program. This should include such techniques as scouting closely for mite activity early in the season, keeping bases of vines and soil surface clean and free of mite-harboring vegetation and initiating selective miticide applications early to break the life cycle and prevent rapid mite population increase.

24(c) Registrant:
Gowan Company
PO Box 5569
Yuma, AZ 85366

Revised 2/12/98